



American Association for Bronchology and Interventional Pulmonology (AABIP) Statement on the Use of Bronchoscopy and Respiratory Specimen Collection in Patients with Suspected or Confirmed COVID-19 Infection

Momen M. Wahidi,* Carla Lamb, MD,* MD, MBA; Septimiu Murgu, MD; Ali Musani, MD; Samira Shojaee, MD; Ashutosh Sachdeva, MD; Fabien Maldonado, MD; Kamran Mahmood, MD; Matthew Kinsey, MD; Sonali Sethi, MD; Amit Mahajan, MD; Adnan Majid, MD; Colleen Keyes, MD; Abdul Hamid Alraiyes, MD; Arthur Sung, MD; David Hsia, MD and George Eapen, MD.

Considering the global spread of COVID-19 infection and the increased number of confirmed COVID-19 cases across the United States, the AABIP is issuing this statement on the safe and effective use of bronchoscopy in patients with suspected or confirmed COVID-19 infection. The main purpose of this statement is to ensure the safety of our patients, health care team and community at large. We are releasing these urgent recommendations to guide clinicians around the world with the understanding that new information may subsequently modify or impact these current recommendations. We will strive to update this statement as needed in a timely fashion. This document is based on the latest Centers for Disease Control (CDC) recommendations and expert consensus of the AABIP COVID-19 Task Force.

General Recommendations for collection of respiratory specimen collection for suspected COVID – 19: (1-4)

- **Collection of upper respiratory samples via nasopharyngeal and oropharyngeal swabs is the primary and preferred method for diagnosis.**
- Respiratory specimen collection is recommended in suspected COVID-19 regardless of time of onset of symptoms.
- **Induced Sputum Collection is NOT recommended.**
- **Because it is an aerosol generating procedure that poses substantial risk to patients and staff, bronchoscopy should have an extremely limited role in diagnosis of COVID-19 and only be considered in intubated patients if upper respiratory samples are negative and other diagnosis is considered that would significantly change clinical management.**
- Alternative respiratory specimen collection in the intubated patient can include tracheal aspirates and non-bronchoscopic alveolar lavage (N-BAL).
- If bronchoscopy is being performed for COVID 19 sample collection, a minimum of 2- 3 ml of specimen into a sterile, leak proof container for specimen collection is recommended. (4)



- Only essential personnel should be present when performing any specimen collection.
- Alert lab personnel regarding COVID-19 specimen processing and testing.

Additional Considerations for Respiratory Evaluation:

- Constellation of fever, respiratory symptoms and radiographic evidence of ground glass opacities and pneumonitis should raise clinical suspicion of COVID-19. (5-6) Patients demonstrating such symptoms or findings should be queried about personal history of recent travel to any country with a CDC Level 2 or higher travel warning (currently China, Iran, South Korea, Japan and most of Europe), contact with a confirmed COVID-19 person or contact with others with such travel history.
- Clinicians should consider the local prevalence of COVID-19 cases when evaluating the clinical risk for COVID-19 infection, understanding that a travel or exposure history will become increasingly ineffective in identifying patients at risk for infection.
- Guidelines for respiratory and contact isolation should be followed in all known or suspected cases of COVID-19 infections.
- Evaluate for influenza and respiratory syncytial virus as well as other respiratory pathogens and additional diagnoses as clinically indicated.
- **For all suspected COVID-19 cases notify internal institutional infection control personnel and state or local public health department.**

General Personnel Preparation if Bronchoscopy is needed in patients with suspected or confirmed COVID-19 infection:

- Place patient in Airborne Infection Isolation Room (AIIR) negative pressure room isolation
- All personnel should wear a **powered, air-purifying respirator** (PAPR) or N95 mask and eye protection
- All personnel should wear standard Personal Protective Equipment (PPE) which includes gown, gloves, respiratory protection, and eye protection
- Follow CDC instructions for proper donning and doffing of all protective equipment and disposable devices. <https://www.cdc.gov/hai/prevent/ppe.html>
- Disposable bronchoscopes should be used first line when available
- Follow standard disinfection protocol of durable re-usable video monitors
- Follow standard High-Level Disinfection for re-usable bronchoscopes
- Limit to essential medical personnel during the procedure and specimen collection.



General Precautions for performing non-urgent bronchoscopy among patients **WITHOUT** suspected COVID-19 infection:

- All patients presenting for previously scheduled bronchoscopy should be asked about their recent travel history prior to entering the bronchoscopy suite. Bronchoscopy should be postponed if the patient has a history of recent travel to any country with a CDC Level 2 or higher travel warning (currently China, Iran, South Korea, Japan and most of Europe).
- All patients should be asked about any fever or ongoing infectious or respiratory symptoms prior to bronchoscopy. Procedures should be postponed if possible until such symptoms have resolved or testing (if available) is negative. If procedures cannot be postponed as determined by the clinical indication, the procedure should be performed using the precautions as outlined above for bronchoscopy in suspected COVID-19 infection.
- In communities with high prevalence of COVID-19 infections, even for routine bronchoscopies in asymptomatic patients, proper isolation precautions should be adhered to while also limiting the number of personnel to essential personnel present in either the bronchoscopy suite or operating room suite with negative pressure room settings or designated isolation room (AIIR).
- **As the rate of community transmission increase, there is increasing concern that asymptomatic patients may present from the community without any pertinent travel or COVID 19 contact history, but may harbor an occult COVID-19 infection. Such concerns are primarily valid in communities with high prevalence of COVID-19 infection and where community transmission has occurred (as opposed to known exposure to travelers). In these communities, the health care team should perform all bronchoscopies wearing appropriate PPE including N95 respirators and face shields. Please note that as testing changes and becomes more widely available, these recommendations will likely change to include routine testing prior to bronchoscopy.**
- Bronchoscopists should consult their local infection control team on when to activate the policy on wearing N95 masks and eye protection routinely for all bronchoscopy procedures in their institution based on local prevalence and community spread of COVID-19 infections.
- This decision must be carefully balanced against the unnecessary use and potential exhaustion of N95 resources. In the case of expected or known N95 respirator shortage, the CDC has advised a strategy of limited re-use of N95 respirators for patients with COVID-19. While not typically within current U.S. standards of care such measures may be considered during crises and epidemics

<https://www.cdc.gov/coronavirus/2019-ncov/hcp/checklist-n95-strategy-h.pdf>

<https://www.cdc.gov/niosh/topics/hcwcontrols/recommendedguidanceextuse.html>



A Call to Postpone non-urgent Bronchoscopy Procedures:

- In order to reduce community spread of COVID-19 infections and preserve healthcare work force and hospital resources, the AABIP is recommending postponing non-urgent bronchoscopy procedures until at least May 2020.
- This is consistent with the recommendations of the American College of Surgeons and the U.S. Surgeon General (<https://www.facs.org/about-acsc/covid-19/information-for-surgeons/triage>)
- We fully recognize that while we prepare to care for our COVID-19 patients, we must also care for other patients, many of whom may have urgent needs. The medical necessity of any bronchoscopy procedure should be judiciously evaluated by the bronchoscopist with careful assessment of the risk of delaying the procedure
- Below is a table that may provide general guidance to bronchoscopists on procedure urgency with the recognition that this is not an exhaustive list nor can it be generalized to every patient

Emergent Bronchoscopy	Urgent Bronchoscopy	Non Urgent Bronchoscopy
Severe or moderate symptomatic Tracheal or Bronchial Stenosis	Lung mass suspicious for cancer	Mild tracheal or bronchial stenosis
Symptomatic central airway obstruction (endotracheal or endobronchial mass or mucus plug)	Mediastinal or hilar adenopathy suspicious for cancer	Clearance of mucus
Massive hemoptysis	Whole lung lavage	High suspicion of sarcoidosis with no immediate need to start therapy
Migrated stent	Foreign object aspiration	Chronic interstitial lung disease
	Mild to moderate hemoptysis	Detection of chronic infection (MAI)
	Suspected pulmonary infection in immunocompromised patients	Chronic cough
		Tracheobronchomalacia evaluation
		Bronchial thermoplasty
		Bronchoscopic lung volume reduction



Indications for Bronchoscopy in patients with suspected or confirmed COVID-19 infections:

- Bronchoscopy is relatively CONTRAINDICATED in patients with suspected and confirmed COVID-19 infections. The only role for bronchoscopy would be when less invasive testing to confirm COVID-19 are inconclusive, suspicion for an alternative diagnosis that would impact clinical management is suspected, or an urgent life-saving intervention as cited below.
- Bronchoscopy for any non-urgent reason should be postponed until after full recovery and the patient is declared free of infection.
- If immediate testing is not available, bronchoscopy should be deferred if possible.
- Bronchoscopy (Flexible and Rigid) for urgent/emergent reasons should be considered only if a lifesaving bronchoscopic intervention is deemed necessary or significant alteration in clinical prognosis is expected with delay.

Information contained in this document will be updated regularly as new information becomes available. For the latest version, please visit <https://aabronchology.org/>

References

1. CDC website <https://www.cdc.gov/coronavirus/2019-nCov/hcp/clinical-criteria.html>
2. Namendys-Silva. Respiratory support for patients with COVID-19 infection. Lancet Respiratory Medicine Epub March 2020
3. Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. China CDC Weekly 2020
4. World Health Organization Laboratory testing for 2019 novel Coronavirus (2019-nCov) in suspected human cases. Interim Guidance 17 Jan 2020
5. Ai T, Yang Z, Hou H., et al. Correlation of chest CT and RT-PCR testing in Coronavirus Disease 2019 (COVID-19) in China: A Report of 1014 Cases online RSNA.org Feb 26, 2020
6. Kanne J, Little B, Chung J, et al. Essentials for Radiologists on COVID-19: An Update online RSNA.org Feb 27, 2020

*Co-first authors.